

VII - CLAIMING

What is claimed is:

1. The procedure for cloning human β A precursor protein gene (human APP gene) based on the reverse transcription (RT) and the polymerase chain reaction (PCR) using the synthesized oligonucleotides (a) for RT, and (b) and (c) for PCR respectively, comprising:
 - Isolation of APP-mRNA.
 - RT reaction using the synthesized oligonucleotide (a) with the following sequence: 5' GTTACAGCACAG 3' (a).
 - RT reaction conditions: 90°C for 2 minutes; 0°C for 1 minute; 25°C for 10 minutes; 42°C for 45 minutes.
 - PCR reaction using the synthesized oligonucleotides (b) and (c) with the following sequences:
5' ATGCTGCCCCGGTTTGGC 3' (b).
5' CTAGTTCTGCATCTGCTCA 3' (c).
 - PCR reaction conditions: Denaturation at 94°C for 1 minutes; annealing at 55°C for 2 minutes; elongation at 72°C for 3 minutes each cycle, for 35 cycles.
2. The procedure for the construction of expression plasmids using the pFastBacTM HTb and the pBlueBacHis2 A transfer vectors for the purpose of obtaining human APP in insect cells, comprising:
 - 2.1. Using the pFastBacTM HTb vector:
 - Digesting the pFastBacTM HTb vector with XbaI and HindIII followed by dephosphorylation with calf intestinal alkaline phosphatase.
 - Digesting the vectors (1) pCR^R II/APP₇₅₁-cDNA and (2) pCR^R II/APP₇₇₀-cDNA with XbaI and HindIII and isolating the resulting fragments containing the cDNA coding sequences of APP, APP₇₅₁-cDNA and APP₇₇₀-cDNA.
 - Ligating the APP₇₅₁-cDNA and APP₇₇₀-cDNA fragments to the pFastBacTM HTb vectors and introducing the ligation products in INV α F⁺ E. Coli strain.
 - Screening for inserts based on the presence of white colonies, as a result of which

the vectors (3) pFastBacTM HTb /APP₇₅₁-cDNA and (4) pFastBacTM HTb /APP₇₇₀-cDNA are selected.

- Introducing the vectors (3) and (4) in DH10BacTM E. Coli competent cells.

- Screening for recombinant bacmids in DH10BacTM E. Coli using blue-white color selection, then verifying the presence of APP-cDNA's inserts in the recombinant bacmids by PCR amplification using the M13 forward (-40) and M13 reverse primers. As a result, the recombinant bacmids (5) for vectors (3) in DH10BacTM E. Coli and (6) for vector (4) in DH10BacTM E. Coli respectively are selected.

2.2. Using the pBlueBacHis2 A vector:

- Digesting the pBlueBacHis2 A vector with NcoI and HindIII followed by dephosphorylation with calf intestinal phosphatase.

- Digesting the vectors (3) pFastBacTM HTb/APP₇₅₁-cDNA and (4) pFastBacTM HTb/APP₇₇₀-cDNA with NcoI and HindIII and isolating the resulting fragments containing the cDNA coding sequences of APP, APP₇₅₁-cDNA and APP₇₇₀-cDNA.

- Ligating the APP₇₅₁-cDNA and APP₇₇₀-cDNA fragments to the pBlueBacHis2 A vectors and introducing the ligation products in INVαF' E. Coli strain.

- Screening for inserts using blue-white color selection, as a result of which the vectors (7) pBlueBacHis2 A/APP₇₅₁-cDNA and (8) pBlueBacHis2 A/APP₇₇₀-cDNA are selected.

3. The procedure for the construction of expression plasmids using the pET-28a (+) transfer vector for the purpose of obtaining human APP in bacteria, comprising:

- Digesting the pET-28a (+) vector with SalI and HindIII followed by dephosphorylation with calf intestinal alkaline phosphatase.

- Digesting the vector (3) pFastBacTM HTb/APP₇₅₁-cDNA and (4) pFastBacTM HTb/APP₇₇₀-cDNA with SalI and HindIII and isolating the resulting fragments containing the cDNA coding sequences of APP, APP₇₅₁-cDNA and APP₇₇₀-cDNA.

- Ligating the APP₇₅₁-cDNA and APP₇₇₀-cDNA fragments to the pET-28a (+) vectors and introducing the ligation products in INVαF' E. Coli strain.

- Screening for inserts based on the presence of white colonies, as a result of which the vectors (9) pET-28a (+)/APP₇₅₁-cDNA and (10) pET-28a (+)/APP₇₇₀-cDNA are selected.

VIII - REFERENCES

- 1- Glenner, G.G. and Wong, C.W.. Alzheimer's disease: Initial report of the purification and
5 characterizaion of a novel cerebrovascular amyloid protein. Biochem. Biophys.
Res.Comm. **120** (1984) 885-890.
- 2- Masters, C.L.; Simms, G.; Weinman, N.A.; Multhaup, G.; McDonalld, B.L.; Beyreuther,
K.. Amyloid plaque core protein in Alzheimer disease and Down syndrome. Proc. Natl.
10 Acad. Sci. USA. **82** (1985) 4245-4249.
- 3- Goldgaber, D.; Lerman, M.I.; McBride, O.W.; Saffiotti, U.; Gajdusek, D.C..
Characterization and chromosomal localization of a cDNA encoding brain amyloid of
Alzheimer's disease. Science **235** (1987) 877-880.
15
- 4- Tanzi, R.E.; Gusella, J.F.; Watkins, P.C.; Bruns, G.A.P.; George-Hyslop, P. H. St.; Van
Keuren, M. L.; Patterson, D.; Pagan, S.; Kurnit, D.M.; Neve, R.L.. Amyloid β protein gene:
cDNA, mRNA distribution, and genetic linkage near the Alzheimer locus. Science **235**
(1987) 880-884.
20
- 5- George-Hyslop, P.H.St.; Tanzi, R.E.; Polinsky, R.J.; et al.. The genetic defect causing
familial Alzheimer's disease map on chromosome 21. Science **235** (1987) 885-890.
- 6- Kang, J.; Lemaire, H.G.; Unterbeck, A.; Salbaum, J.M.; Masters, C.L.; Grzeschik, K.H.;
25 Multhaup, G.; Beyreuther, K.; Muller-Hill, B.. The precursor of Alzheimer's disease
amyloid A4 protein resembles a cell-surface receptor. Nature **325** (1987) 733-736.
(GenBank Accession No. Y00264).
- 7- Ponte, P.; Gonzalez-DeWhitt, P.; Schilling, J.; Miller, J.; Hsu, D.; Greenberg, B.; Davis,
30 K.; Wallace, W.; Lieberburg, I.; Fuller, F.; Cordell, B.. A new A4 amyloid mRNA contains a
domain homologous to serine proteinase inhibitors. Nature **331** (1988) 525-527. (GenBank
Accession No. Y00297).
- 8- Tanzi, R.E.; McClatchey, A.I.; Lamperti, E.D.; Vill-Komaroff, L.; Gusella, J.F.; Neve,
35 R.L.. Protease inhibitor domain encoded by an amyloid protein precursor mRNA associated
with Alzheimer's disease. Nature **331** (1988) 528-530.
- 9- Kitaguchi, N.; Takahashi, Y.; Tokushima, Y.; Shiojiri, S.; Ito, H.. Novel precursor of
Alzheimer's disease amyloid protein shows protease inhibitory activity. Nature **331** (1988)
40 530-532.
- 10- Oltersdorf, T.; Fritz, L.C.; Schenk, D.B.; Lieberburg, I.; Johnson-Wood, K.L.; Beattie,
E.C.; Ward, P.J.; Blacher, R.W.; Dovey, H.F.; Sinha, S.. The secreted form of the
Alzheimer's amyloid precursor protein with the Kunitz domain is protease nexin-II. Nature
45 **341** (1989) 144-147.

- 11- Van Nostrand, W.; Schmaier, A.H.; Farrow, J.S.; Cunningham, D.D.. Protease nexin II (amyloid β -protein precursor): A platelet α -granule protein. Science **248** (1990) 745-748.
- 12- Smith, R.P.; Higuchi, D.A.; Broze Jr, G.J.. Platelet coagulation factor XIa-inhibitor, a form of Alzheimer amyloid precursor protein.. Science **248** (1990) 1126-1128.
- 13- Schubert, D.; Cole, G.; Saitoh, T.; Oltersdorf, T.. Amyloid beta protein precursor is a mitogen. Biochem. Biophys. Res. Commun. **162** (1989) 83-88.
- 14- Schubert, D.; Jin, L.W.; Saitoh, T.; Cole, G.. The regulation of amyloid β protein precursor secretion and its modulatory role in cell adhesion. Neuron **3** (1989) 689-694.
- 15- Yankner, B.A.; Duffy, L.K.; Kirschner, D.A.. Neurotrophic and neurotoxic effects of amyloid β protein: Reversal by tachykinin neuropeptides. Science **250** (1990) 279-282.
- 16- Whitson, J.S.; Selkoe, D.J.; Cotman, C.W.. Amyloid-protein enhances the survival of hippocampal neurons in vitro. Science **243** (1989) 1488-1490.
- 17- Yankner, B.A.; Dawes, L.R.; Fisher, S.; Villa-Komaroff, L.; Oster-Granite, M.L.; Neve, R.L.. Neurotoxicity of a fragment of amyloid precursor associated with Alzheimer's disease. Science **245** (1989) 417-420.
- 18- Weidemann A.; Konig, G.; Bunke, D.; Fischer, P.; Salbaum, J.M.; Masters, C.L.; Beyreuther, K.. Identification, biogenesis, and localization of precursors of Alzheimer's disease A4 amyloid protein. Cell **57** (1989) 115-126.
- 19- Oltersdorf, T.; Ward, P.J.; Henriksson, T.; Beattie, E.C.; Neve, R.; Lieberburg, I.; Fritz, C.. The Alzheimer amyloid precursor protein. Identification of a stable intermediate in the biosynthetic/degradative pathway. J. Biol. Chem. **265** (1990) 4492-4497.
- 20- Knops, J.; Johnson-Wood, K.; Schenk, D.B.; Sinha, S.; Lieberburg, I.; McConlogue, L.. Isolation of baculovirus-derived secreted and Esch, F.S.full-length-amyloid precursor protein. J. Biol. Chem. **265** (1991) 7285-7290.
- 21- Esch, F.S.; Keim, P.S.; Beattie, E.C.; Blacher, R.W.; Culwell, A.R.; Oltersdorf, T.; McClure, D.; Ward, P.J.. Cleavage of amyloid β peptide during constitutive processing of its precursor. Science **248** (1990) 1122-1124.
- 22- Suzuki, N.; Cheung, T.T.; Cai, X.D.; Odaka, A.; Otvos Jr, L.; Eckman, C.; Golde, T.E.; Younkin, S.G.. An increase percentage of long amyloid β protein secreted by familial amyloid β protein precursor (β APP₇₁₇) mutants. Science **264** (1994) 1336-1340.
- 23- Cai, X.D.; Golde, T. E.; Younkin, S.G.. Release of excess amyloid β -protein from a mutant amyloid β -protein precursor. Science **259** (1993) 514-516.

24- Hsiao, K.; Chapman, P.; Nilsen, S.; Eckman, C.; Harigaya, Y.; Younkin, S.; Yang, F.; Cole, G.. Correlative memory deficits, A β elevation, and amyloid plaques in transgenic mice. Science **274** (1996) 99-102.

5 25- Sambrook, J.; Fritsch, E.F.; Maniatis, T.. Extraction, purification and analysis of messenger RNA from eukaryotic cells. In molecular cloning, a laboratory manual, 2nd Ed.; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, New York, 1989; 7.28-7.52.

10 26- Saiki, R.K.; Scharf, S.; Faloona, F.; Mullis, K.B.; Horn, C.T.; Erlich, H.A.; Arnheim, N.. Amplification of β -globine genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. Science **230** (1985) 1350-1354.

15 27- Kawasaki, E.S.; Wang, A.M.. Detection of gene expression. In PCR technology; Erlich, H.A.; Ed. Stockton: New York, 1989, 89-97.